## Treatment Protocols Drowning/Near Drowning - Adult

Drowning/Near Drowning - Adult	<b>Policy #9090</b> A
Adult BLS Standing Orders	
Universal Patient Protocol	
• Ensure patent airway, suction PRN	
• Intervene PRN per Airway Policy	
• Spinal motion restriction PRN per Spinal Motion Restriction Policy	
<ul> <li>Monitor O2 saturation, aggressively treat hypoxia or increased WOB</li> </ul>	
<ul> <li>Give oxygen and/or ventilate PRN</li> </ul>	
<ul> <li>Keep patient warm</li> </ul>	
<ul> <li>All patients should be evaluated in the hospital, BHP</li> </ul>	
<ul> <li>Blood glucose PRN</li> </ul>	
<ul> <li>Refer to Poisoning and Trauma Policies as needed</li> </ul>	
<ul> <li>Capnography</li> </ul>	
• CPAP per NIPPV Policy	
Adult LALS Standing Order Protocol	
Establish IV PRN	
• Capnography	
Wheezing	
• Albuterol 2.5 mg nebulized for wheezing MR x 1	
Hypotension	
Refer to Shock Protocol	
Adult ALS Standing Order Protocol	
Monitor EKG	
• IV/IO PRN	
Capnography	
• ETT prn per Airway Policy	
• CPAP per NIPPV Policy	
Wheezing	
• Albuterol 2.5 mg nebulized for wheezing MR x, 1 MR <b>BH</b>	
Hypotension	
• NS 500-1,000 mL IV MR x 1 to a max of 2,000 mL to maintain a SBP of $\geq$ 90 mmH	g
Refer to Shock Protocol	
Nausea and/or Vomiting	
• Ondansetron 4 mg IV/IO/IM/ODT PRN, MR x1, total 8 mg	
Adult Base Hospital Orders	
• <b>BH</b> – Repeat albuterol 2.5 mg PRN wheezing	
Notes:	

• All patients should be evaluated in the hospital. Consult with **BHP** if patient is desiring AMA

APPROVED: SIGNATURE ON FILE – 07/01/25 Katherine Staats, M.D. FACEP EMS Medical Director