Treatment Protocols Drowning/Near Drowning – Pediatric

Pediatric BLS Standing Orders Universal Patient Protocol • Ensure patent airway, suction PRN • Intervene PRN per Airway Policy • • Spinal motion restriction PRN per Spinal Motion Restriction Policy Monitor O2 saturation, aggressively treat hypoxia or increased WOB ٠ • Give oxygen and/or ventilate PRN • Keep patient warm All patients should be evaluated in the hospital, **BHP** ٠ Blood glucose PRN • Refer to **Poisoning** and **Trauma Policies** as needed • Capnography • **Pediatric LALS Standing Orders** Establish IV PRN • Capnography • Wheezing • Albuterol – per dosing chart via nebulizer MR x1 BH Hypotension • Follow Shock Protocol **Pediatric ALS Standing Orders** • Monitor EKG IV/IO PRN • Capnography • **Refer to Airway Policy** Wheezing Albuterol – per dosing chart via nebulizer MR x1 • Hypotension • 10-20 mL/kg NS IV bolus; titrated to age-appropriate systolic BP MR x1 • Use Shock Protocol for persistent hypotension Nausea and/or Vomiting • Ondansetron 0.1 mg/kg, max 4 mg - IV/IO/IM/ODT PRN MR x1 **Pediatric Base Hospital Orders BH** – Repeat albuterol weight based nebulized **BH** – Repeat NS bolus IV/IO PRN hypotension •

Notes:

• All patients should be evaluated in the hospital. Consult with **BHP** if requested for AMA.

Treatment Protocols <u>*Drowning/Near Drowning – Pediatric*</u> APPROVED:

SIGNATURE ON FILE - 07/01/25

Katherine Staats, M.D. FACEP

EMS Medical Director