

Treatment Protocols**Date: 07/01/2023*****Behavioral Emergencies - Adult*****Policy #9050A****Adult BLS Standing Orders**

- **Universal Patient Protocol**
- **Oxygen** – PRN
- Blood glucose test
- Attempt to determine illness or mechanism, consider **Spinal Motion Restriction, Altered Mental Status, and Trauma** policies
- Restrain only if necessary, refer to **Patient Restraint Policy**
- TASER® probes are to be left in place, only remove if affecting airway or lifesaving treatment.
- Consider law enforcement support
- Consider 5150 evaluation
- Attempt verbal de-escalation primarily. If unsuccessful, consider physical and chemical restraints as needed and able based on patient and provider safety

Adult LALS Standing Order Protocol

- Establish IV PRN

Adult ALS Standing Order Protocol

- Monitor EKG
- Establish IV/IO PRN
- Capnography PRN (required if patient receives midazolam)
- Obtain 12 Lead EKG if patient was tased, patient is unstable, or source of behavior change is unknown

Patients Exhibiting Severe Agitation

Midazolam 5 mg IV/IN/IM, MR x1 q2min to a max dose of 10 mg. MR BH

For patients that are elderly, small statured, have other medications, intoxicants, or medical sources for their agitation, consider administering midazolam in 1 mg or smaller increments, waiting at least 2 minutes between doses.

- Do not exceed 1.0 mL per nostril per dose for adult patients

Adult Base Hospital Orders**Patients Exhibiting Severe Agitation**

- **BH** – Further midazolam doses up to 10 mg, MR x 2. (30 mg total)
- Monitor and anticipate respiratory depression with larger doses

Treatment Protocols**Date: 07/01/2023*****Behavioral Emergencies - Adult*****Policy #9050A****Notes**

- IM is preferred route of administration due to risk of injury to patient or EMS personnel.
- Consult with **BH** prior to releasing patient in custody of law enforcement, or other legally responsible party. If a patient has received medication from EMS, the patient is NOT appropriate for release to custody.
- **If a patient received midazolam for agitation, they are not appropriate to AMA.**
- Be aware if patients are intoxicated, midazolam may suppress their respiratory drive, or cause total apnea and the patient may require respiratory support.
- It is always critical with behavioral emergencies to determine what potential medical cause could be contributing to the patient's behavior. Ensure clinical consideration (and evaluation) for intracranial bleeding or injury, low or high glucose, infection, medication overdoses, or other causes of altered mental status.

APPROVED:

Signature on File

Katherine Staats, M.D. FACEP

EMS Medical Director