

Treatment Protocols

Date: 02/01/2021

Universal Protocol

Policy #9000

| Medical | Trauma |
|--|--|
| Adult and Pediatric BLS Standing Orders | |
| <ul style="list-style-type: none"> • Evaluate scene safety • Assess scene needs and resources and activate further assistance as necessary • Personal protective equipment • Assess, establish and maintain airway per Airway Policy • Evaluate breathing and circulation • Assess chief complaint • Perform focused assessment and physical exam • Glucose Measurement PRN • Vital signs: <ul style="list-style-type: none"> ○ HR, BP, RR, Pulse Oximetry, Pain assessment ○ Reassess frequently • Assess need for ALS Rendezvous • BLS Treatment Protocols • Transport Determination | <ul style="list-style-type: none"> • Evaluate scene safety • Assess scene needs and resources and activate further assistance as necessary • Personal protective equipment • Assess, establish and maintain airway per Airway Policy • Establish Spinal Motion Restriction per policy • Evaluate breathing and circulation • Remove patient’s clothing to expose and identify injuries • Ensure patient warmth – recover patient after clothing removal to maintain core body temperature • Perform focused assessment and physical exam • Glucose Measurement PRN • Vital signs: <ul style="list-style-type: none"> ○ HR, BP, RR, Pulse Oximetry, Pain assessment ○ Reassess frequently • Assess need for ALS Rendezvous • BLS Treatment Protocol • Transport Determination |
| LALS Standing Order Protocols | |
| <ul style="list-style-type: none"> • Establish IV PRN • LALS Treatment Protocols PRN | <ul style="list-style-type: none"> • Establish IV PRN • LALS Treatment Protocols PRN |
| Adult ALS Standing Order Protocols | |
| <ul style="list-style-type: none"> • 12 lead PRN • Establish IO PRN • ETCO2 monitoring PRN • ALS Treatment Protocols • Transport Determination | <ul style="list-style-type: none"> • 12 lead PRN • Establish IO PRN • ETCO2 monitoring PRN • ALS Treatment Protocols • Transport Determination |
| Note | |
| <ul style="list-style-type: none"> • Pediatric patients fitting on the pediatric measurement tape (generally this corresponds to younger than 15 years old) • Utilize pediatric measurement tape when applicable for pediatric patients | |

APPROVED:

Signature on File

Katherine Staats, M.D
EMS Medical Director