

**Treatment Protocols****Date: 02/01/2021*****Hemodialysis Related Emergency/Suspected Hyperkalemia*****Policy #9130P****Pediatric BLS Standing Orders**

- **Universal Patient Protocol**
- Give oxygen and/or ventilate PRN per **Airway Policy**
- Continuously monitor O2 saturation, and blood pressure
- Glucose testing PRN
- For bleeding dialysis fistulas or shunts, refer to **Hemorrhage Control Protocol**
- For bradycardia or tachycardia due to suspected electrolyte abnormalities, go to **Dysrhythmia Protocols**

**Pediatric LALS Standing Orders**

- Place IV PRN
- Provide 10-20 ml/kg IV fluid for hypotension PRN

**Pediatric ALS Standing Orders**

- Administer continuous ECG, blood pressure, pulse oximetry, and capnography monitoring PRN
- Place IO PRN
- Obtain 12 lead EKG PRN
- Administer 10-20 ml/kg IO fluid for hypotension PRN without evidence of fluid overload

**Pediatric Base Hospital Orders**

- **BHP** – Consult for treatment of arrhythmias, volume overload, or other suspected result of missed hemodialysis, and/or associated kidney failure

APPROVED:

Signature on FileKatherine Staats, M.D.  
EMS Medical Director